





Aquaculture planning, development in Brazilian federal waters

1 July 2015

By Rodrigo Roubach, Ph.D. , Antônio Ostrensky, Ph.D. , Eduardo Pickler Schulter and Dr. Guilherme Wolff Bueno

Tilapia production will lead the way



A key strategy of Brazil's More Fishing and Aquaculture plan is the development of aquaculture in federal waters. A variety of freshwater fish are raised in the floating cages.

Aquaculture in Brazilian federal waters is worth around 3.0 billion Brazilian reals (U.S. \$987.4 million) in business and provides a million direct and indirect jobs, and feed production around 900 million metric tons (MMT) annually. According to the Brazilian Ministry of Fisheries Aquaculture Institute of Geography and Statistics, this agribusiness produced around 476,000 MT in 2013, representing a growth rate of 56 percent in the past 12 years. Thus, the aquaculture industry is consolidating as the main source for fisheries production, a trend observed in other countries of the world, like China, Norway and Chile.

The Brazilian Institute of Geography and Statistics reported that among the species produced in the country, freshwater fish such as tilapia, *Oreochromis niloticus*, accounted for 43.0 percent of the Brazilian aquaculture production in 2013. Farming of tambaqui, *Colossoma macropomum*, reflected 22.6 percent, and the hybrid "tambacu" (female *C. macropomum* x male *Piaractus mesopotamicus*) made up 15.4 percent of the production. Still, other species have potential to be produced on a large scale, mainly due to their wide acceptance by consumers. These include arapaima, *Arapaima gigas*; surubins, the spotted catfish *Pseudoplatystoma corruscans*; and cachara, *Pseudoplatystoma fasciatum*. However, these species still require further research and development of specific technologies for production on a commercial scale, which happened earlier for tilapia.

Aquaculture In Federal Waters

The current annual world fish production is estimated at 158 MMT, and according to United Nations Food and Agriculture Organization projections for 2030, it is expected to increase around 100 MMT/year. With an estimated potential annual yield of 20 MMT, Brazil could help to meet this demand.

Given these projections, Brazil's federal government – with the participation of the production sector and civil organizations – drafted the sustainable development plan entitled "More Fishing and Aquaculture" in 2008. The objective of the plan was to promote the sustainable development of fisheries and aquaculture by linking all those involved with the sectors, thus consolidating sustainable state development policies addressing social inclusion, security and food sovereignty in Brazil.

The use of reservoirs for multiple uses, including aquaculture, in Brazil was established by the National Water Resources Policy act in 1997. This legal framework was complemented by federal directives regulating the use of Brazilian waters and public spaces for the practice of aquaculture. Up to 1 percent of the area of federal waters is available for aquaculture, which corresponds to at least 55,000 ha in freshwater reservoirs.

Reservoir studies

A key strategy of the More Fishing and Aquaculture plan is the development of aquaculture in federal waters. Therefore, the Ministry of Fisheries and Aquaculture has invested in studies for the demarcation of aquaculture parks in many reservoirs, as well as the coastal areas of Santa Catarina and Paraná.

These studies were based on criteria that addressed issues of environmental, social and economic sustainability. These 10 reservoirs have potential for the implementation of 100 aquaculture parks with a total area over 1,920 hectares for fish culture and production capacity estimated at 227,000 MT/year of fish. This number would rise to more than 284,000 MT if the two studies in coastal areas are included (Table 1).

Roubach, Estimated production for demarcated aquaculture parks, Table 1

Area	Reservoir/State	State	Estimated Production (mt/year)	Most Important Species
Reservoir	Furnas	Minas Gerais	35,702	Oreochromis niloticus
Reservoir	Três Marias	Minas Gerais	25,748	Oreochromis niloticus
Reservoir	Ilha Solteira	Mato Grosso do Sul/ São Paulo/ Minas Gerais	50,422	Oreochromis niloticus
Reservoir	Castanhão	Ceará	33,080	Oreochromis niloticus
Reservoir	Itaipu	Itaipu	2,920	Piaractus mesopotamicus
Reservoir	Tucuruí	Pará	8,518	Piaractus brachypomus
Reservoir	Lajeado	Tocantins	21,856	Colossoma macropomum
Reservoir	Serra da Mesa	Goiás	31,356	Oreochromis niloticus
Reservoir	Cana Brava	Goiás	10,140	Oreochromis niloticus
Reservoir	Manso	Mato Grosso	7,192	Piaractus mesopotamicus
Coastal	Santa Catarina	Santa Catarina	306	Crassostrea gigas, Perna perna, Nodipecten nodosus
Coastal	Paraná	Paraná	57,538	Crassostrea rizophora
Total	-	_	284,778	-

Source: SINAU/MPA, 2015

Currently, studies are being finalized and/or aquaculture parks are being implemented in 36 reservoirs: five in the Midwest, eight in the Northeast, five in the North, eight in the Southeast and 10 in the South. This development represents at least 210 parks. Of these, about 100 will be allocated to aquaculture on a family scale.

Furthermore, in recent years, studies have been conducted to set up additional marine aquaculture parks in several states where parks for the production of fish, oysters, scallops, mussels and algae are already occupied by producers/entrepreneurs. The parks already identified may contribute increases of 59 percent over the current national aquaculture production and 37 percent in the present national fisheries production.

Carrying capacity

The granting of permission to use freshwater bodies for aquaculture projects in Brazil is carried out by the National Water Agency, which sets the maximum permissible value of fish in a given reservoir by calculating the carrying capacity using the hydrodynamic model developed by Peter J. Dillon and Frank H. Rigler in 1974. However, new research and pilot projects are being developed by University of Brasilia with Guelph University under a collaborative project involving the Brazilian Ministry of Science and Technology's National Council for Scientific and Technological Development, and International Science and Technology Partnerships Canada to improve the methodology and input data through the integration of a nutritional bioenergetics model described by C. Young Cho and Dominique P. Bureau in 1998.

The hydrodynamic model projects waste dispersion simulations for the use of commercial diets with different total phosphorus percentage. This model allows more compartmentalized analysis of aquaculture parks, assigning technical, environmental and nutritional aspects for each region and increasing the security and reliability of estimates of environmental carrying capacity.

To enable the safe and environmentally responsible implementation of aquaculture in federal waters, mechanisms will be established to support supervised management of fish culture areas and define maximum productivity indicators, safe phosphorus levels in feed and distances between culture areas to allow appropriate dilution of waste in the environment.

Brazil's great climate, considerable species diversity and abundance of water resources have combined to support aquaculture expansion in the last 10 years.

Aquaculture potential, challenges

Brazil is a country with great climatic diversity, great species diversity and an abundance of water resources for the development of aquaculture in federal waters. Aquaculture activity in lakes and reservoirs has expanded in the last 10 years, and hence the volume of projects and farms registered by the Ministry of Fisheries and Aquaculture.

However, despite this potential, there are a number of challenges to overcome for consolidation of the Brazilian aquaculture program. The main challenges are the lack of standardization of environmental licensing procedures across the states of the federation, bureaucracy and the limited preparation of environmental agencies to review projects and issue environmental licenses.

Although tilapia is the primary species farmed in Brazil, tambaqui (a), pacu (b), cachara (c) and pirarucu (d) are also cultured to meet consumer demand.

These problems have increased the risk of investment in aquaculture, especially when factors such as global climate change, localized water scarcity and depletion of reservoirs, eutrophication and animal health problems are also considered.

An important point to examine is the absence of specific biosafety and health policies for aquaculture in federal waters, since the aquaculture parks program groups together entrepreneurs, family farmers and government projects with different objectives and different degrees of knowledge and concern. Thus, there is a need to implement management committees and programs for monitoring and control of the reservoirs and the Brazilian coast.

Finally, integrated and sustainable development of the production chain in these regions – with tax exemption in the acquisition of supplies and equipment – is needed, along with coordination across the production chain to minimize the economic, social and environmental risks of these projects in Brazilian federal waters.

(Editor's Note: This article was originally published in the July/August 2015 print edition of the Global Aquaculture Advocate.*)*

Authors



RODRIGO ROUBACH, PH.D.

Ministry of Fisheries and Aquaculture SBS Quadra 02, Lote 10 Bloco J, Edificio Carlton Tower Brasília, Distrito Federal, Brazil

rodrigo.roubach@mpa.gov.br (mailto:rodrigo.roubach@mpa.gov.br)



ANTÔNIO OSTRENSKY, PH.D.

Department of Animal Science Federal University of Paraná Curitiba, Paraná, Brazil



EDUARDO PICKLER SCHULTER

EPS Aquaculture and Environment



DR. GUILHERME WOLFF BUENO

Department of Animal Science University of Brasília Campus Universitário Darcy Ribeiro Brasília, Distrito Federal, Brazil Copyright © 2023 Global Seafood Alliance

All rights reserved.