



ALLIANCE™

[.https://debug.globalseafood.org](https://debug.globalseafood.org)Health &  
Welfare

# Examining the protozoan Perkinsus sp. in Mexican Pacific oyster aquaculture

6 May 2016

By Cesar M. Escobedo-Bonilla, Ph.D. and Lizeth C. Villanueva-Fonseca, M.C.

## Study determines prevalence, infection intensity in Northern Sinaloa

The Pacific oyster (*Crassostrea gigas*), a mollusk native to Japan, is a well-regarded and appreciated species for aquaculture due to its fast growth rate and ability to withstand wide salinity variations. At present, this species is cultured worldwide in more than 20 countries. In Mexico, this species was first introduced in San Quintin Bay, Baja California in 1973. Since then, its farming has spread throughout northwest Mexico.



The Pacific oyster is an important aquaculture species that is farmed in many regions around the world.

## Oyster pathogen study

In Mexico, the presence of pathogens affecting this species has been recorded since 1962, when the protozoan *Perkinsus marinus* was found in the American oyster (*Crassostrea virginica*) in Laguna Madre, Tamaulipas. In the late 1990s, this pathogen was reported in the eastern states of Tabasco and Veracruz. The Pacific coast of Mexico was considered free of this pathogen, but mortality records in this area led to monitoring the health status of the farmed oyster species in the region.

In 1998, high mortalities were first recorded in farmed oysters along the Gulf of California. The causative agent was not recognized and it was thought that a set of factors such as high water temperature, high stocking density, presence of pollutants along with physiological stress due to reproduction and the presence of pathogens may have been involved. In 2008, the presence of *Perkinsus* sp. was reported in the states of Nayarit, Sonora and Sinaloa in the oyster species *C. cortziensis*, *C. gigas* and *Saccostrea palmula*, respectively. The origin of this protozoan in northwest Mexico is unknown, but it could be associated to the introduction of *C. virginica* from the East Coast of the United States or oysters from the Gulf of Mexico.

The aim of our study was to determine the prevalence of *Perkinsus* sp. in cultured *C. gigas* in the estuary called “La Pitahaya” in northern Sinaloa and to determine the intensity of infection during an annual culture cycle using two detection methods. We are gratefully indebted to Dr. Nancy Stokes, from the Virginia Institute of Marine Science, William & Mary University, Virginia, USA, for the kind donation of *Perkinsus marinus* samples.

The study area was located in “La Pitahaya” estuary, northern Sinaloa, between the coordinates 25 degrees 18’ – 25 degrees 24’ N and 108 degrees 32’ – 108 degrees 45’ W (Fig. 1). Oysters (n = 6000) (size 3-4 mm) from a hatchery in Sonora (CREMES) were used. The oysters were stocked in March at two densities: 28 and 42 oysters per tray (50 x 50 x 10 cm). The long-line culture system was used and the culture was done for 13 months. Every fortnight, water temperature and salinity were monitored, trays were cleaned and mortalities were recorded. Also, oysters (n=50) were sampled for total weight (g) in order to determine the specific growth rate.




Fig. 1: Geographic location of *Crassostrea gigas* culture site in the estuary “La Pitahaya,” Northern Sinaloa, Mexico.

## **Perkinsus** sp. prevalence, infection intensity

For the detection of *Perkinsus* sp., each month 30 oysters were collected at random and they were processed for Thioglycollate Fluid Medium (TFM) assay. Briefly, tissues from gills and/or mantle from a single individual were dissected and placed into a 15 ml Falcon tube containing 9.5 ml sterile TFM added with antibiotics/antimycotics. Incubation was done at 22 to 25 degrees C for 4 to 7 days in the dark. After incubation, the fragments of tissues were collected and minced with a sterile blade on a

glass slide. Tissues were stained with Lugol-iodine solution, covered with a coverslip, incubated at room temperature for 10 min and examined by light microscopy (Leica DM750, Germany) at 100 X and 400 X, respectively.

For PCR detection, gill and mantle tissues were used for DNA extraction (DNAzol, Molecular Research Center, Inc. Cincinnati OH, USA), following the manufacturer's instructions. Primers PerkITS85 (5'-CCG CTT TGT TTG GAT CCC-3') and PerkITS750 (5'-ACA TCA GGC CTT CTA ATG ATG-3') that amplify a fragment (703 base pairs) of the rRNA internal transcribed spacer (ITS) from *Perkinsus* were used (OIE, 2009). A positive DNA control of *Crassostrea ariakensis* infected with *Perkinsus marinus* was used. PCR products were resolved in 1 percent agarose gel electrophoresis stained with ethidium bromide (1 µg/ml).

We determined the monthly prevalence of *Perkinsus* in the culture, as well as the intensity of infection, which was calculated with hyphospore counts using the relative scale of Mackin.





Fig. 3: Detection of *Perkinsus* sp. by the Thioglycollate Fluid Medium (TFM) assay. a) oyster with signs of *Perkinsus* disease; b) mantle tissue cultured in TFM showing no *Perkinsus* hypnospores (150X); c, d) mantle tissues showing *Perkinsus* hypnospores in a level of infection 2, according to the Mackin scale (150 X).

## Results

Oysters showed a specific growth rate of  $2.52 \pm 1.00$  g at density of 28 animals/tray and  $2.54 \pm 1.05$  g at density of 42 animals/tray, respectively. Water temperature was the most influencing factor in growth, since the lowest growth rates were obtained either during the months with highest temperature records in the year ( $29 - 32^\circ\text{C}$ ) (June to October) or during the months with the lowest temperatures ( $\leq 17$  degreesC) (January to April) (Fig. 2). These data showed that temperatures higher than  $27^\circ\text{C}$  or lower than  $22^\circ\text{C}$  have a negative impact in growth. Negative growth rates were recorded in July and October. Here, oysters undergo physiological changes related to sexual maturity and reproduction, which induces weight loss due to gonad reduction.

At the end of the culture, the mean weight of oysters was  $204.07 \pm 18.97$  g and  $207.09 \pm 15.29$  g at densities of 28 and 42 animals/tray, respectively. Survival at the end of culture was 95.9 and 94.3 percent for densities of 28 and 42 oysters/tray, respectively. Mortalities were mostly recorded during the high temperature months (June to October).

Detection of *Perkinsus* infection by TFM first occurred two months after stocking. Prevalence of *Perkinsus* varied between 16.6 and 23.3 percent during the warmest months (August – October), whereas in the coldest months (December to February) it diminished to 20 – 10 percent, respectively. Prevalence of *Perkinsus* was the highest during the last two months of the culture (March to April, 2012) (26.6 and 40 percent, respectively). The intensity of infection by TFM was always between levels 1-2 according to the Mackin scale (Fig. 3).

By PCR, detection of *Perkinsus* occurred until five months after stocking. The prevalence found by PCR was always lower than that obtained by TFM. By PCR, the highest prevalence was found from October to December (10, 6.6 and 13.3 percent, respectively), reducing in January to February, 2012 (6.6 and 0 percent, respectively) and by the end of the culture cycle (March to April, 2012), the prevalence was the highest (16.6 and 26.6 percent, respectively) (Fig. 4).

Fig. 4: Monthly prevalence of *Perkinsus* infection during the 13 months of culture through TFM or PCR diagnosis.

## Perspectives

The presence of *Perkinsus* was recently reported in four coastal areas in Sinaloa, (Cospita, La Reforma, Topolobampo and La Bocanita) in wild populations of oysters *Saccostrea palmula* and *C. corteziensis*. Our study enhances the geographical record of *Perkinsus* sp. in cultured *Crassostrea gigas* in the estuary "La Pitahaya," northern Sinaloa. In our study, prevalence of *Perkinsus* varied in a culture cycle between 40 percent and 6.6 percent, depending on the temperature and the age of the culture. The intensity of infection was level 2, probably due to the relatively low susceptibility of *C. gigas* to *Perkinsus* infection.

## Authors

---



### **CESAR M. ESCOBEDO-BONILLA, PH.D.**

Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional  
Instituto Politécnico Nacional  
Unidad Sinaloa  
Blvd. Juan de Dios Batiz Paredes 250  
Colonia San Joachin, Guasave  
Sinaloa 81101 - México

[cesar\\_escobedomx@yahoo.com](mailto:cesar_escobedomx@yahoo.com) ([mailto:cesar\\_escobedomx@yahoo.com](mailto:cesar_escobedomx@yahoo.com))



### **LIZETH C. VILLANUEVA-FONSECA, M.C.**

Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional  
Instituto Politécnico Nacional  
Unidad Sinaloa  
Blvd. Juan de Dios Batiz Paredes 250  
Colonia San Joachin, Guasave  
Sinaloa 81101 - México

Copyright © 2023 Global Seafood Alliance

All rights reserved.