




(<https://debug.globalseafood.org>).



Innovation &  
Investment

# Experimental aquaponics system in Costa Rica

26 February 2018

By José Valverde , Marco Acosta and Marvin Torres

## Assessing production of channel catfish, freshwater prawns and lettuce



The aquaponic system used in the study included, from left to right: the culture tanks for catfish and prawns; the outlet pipes and the collection tank; the filtration tank with bio balls; and the substrate bed for hydroponic culture of lettuces.

In aquaculture systems, cultured organisms generate waste from their defecation, by the decomposition of non-ingested food and from the contribution of microorganisms that develop around them. This is wastewater rich in nutrients that must leave the system in the form of effluents to avoid contaminating the culture medium. If not removed, some wastes can reach toxic levels for the cultured organisms, so they must be discarded or treated. The treatment of these wastes by conventional methods is usually expensive and complicated.

In an aquaponics system, these wastes are reused as liquid fertilizers to feed the plants through an integration of aquaculture with hydroponics, and a wide variety of plants – growing on different substrates – are able to absorb, through their roots, the nutrients that come from aquatic cultures. Plants act as a natural filter by reducing the concentration of extra nutrients and cleaning the water, which can then be re-integrated and recirculated into the aquatic species component system (recirculating aquaculture system, RAS).

Aquaponics creates a mini-ecosystem between plants and aquatic organisms that benefit and support each other to survive and grow. It has become an ideal alternative because it moves nutrient-rich water from the aquaculture component and uses it as a nutrient for plants in the hydroponics component. In our study, we evaluated the performance of channel catfish and freshwater prawns and three varieties of lettuce – American, Romaine and Rose – in a research aquaponics system.

## Study setup

The study was conducted at the Fishing Nautical Nucleus (NNP) of the National Learning Institute (INA) located in the central canton of the Province of Puntarenas. Carried out under greenhouse conditions, the only water used as liquid fertilizer for the plants came from 1 cubic meter tanks where freshwater prawns (*Macrobrachium rosenbergii*) and channel catfish (*Ictalurus punctatus*) were cultured.

The aquaponics system used in the study included a greenhouse with a translucent plastic roof and walls lined with white anti-aphid mesh. The plastic, square culture tanks (1.15 x 0.95 m, 0.93 m deep) were raised by 1 meter with a wooden platform.

An intensive farming system was used, with the catfish juveniles stocked at a density of 100 per cubic meter and a weight of 0.5 grams, while for the freshwater prawns postlarvae (PL; 0.003 g), two stockings were done one month apart at a density of 100/cubic meters each, for a total stocking of 200 PL per cubic meter. The catfish and prawns were cultured for 85 days with constant aeration and continuous water exchange and fed *ad libitum* with commercial feed for tilapia and marine shrimp, respectively. At the end of the study, the catfish and the prawns were harvested and weighed and returned to the culture system.

After 40 days of culture, the lettuce plants were planted with the recirculation system in operation. In total, 63 plants were planted in an area of 5 square meters for a density of 12.6 lettuces per square meter. The different lettuce varieties were planted interspersed in seven rows of three plants each. The substrate bed was 20 cm deep and the water flowed at 15 cm deep at a flow rate of 200 mL per second.

Various water quality parameters, including temperature, pH and dissolved oxygen, were monitored weekly. Dissolved oxygen levels were maintained at saturation through constant aeration from the continuously operating blower and diffuser hoses.

The plant project lasted for 45 days and the catfish and prawn project for 85 days of culture. The plants were harvested and measured with a tape measure. The aquatic organisms were weighed and measured individually. An evaluation of weekly growth (grams per week), final biomass (grams per



View of the aquaponic system with elevated tanks for the culture of catfish and prawns at the NNP-INA in Puntarenas, Costa Rica.

cubic meter), feeding rate (grams of feed per cubic meter X days) and feed conversion (grams feed per grams of biomass) was made for the catfish and prawns.



Three varieties of lettuce were used in this study, from left to right: American, Romaine and Rose.

## Results

A closed recirculation system was developed and operated where wastewater from the catfish and prawns was removed from the bottom with daily water exchanges. Wastewater went to a collection tank and then to a biological filtration tank with bio balls as a substrate for bacteria.

The nitrifying bacteria convert the ammoniacal compounds from the wastewater to less toxic forms for the catfish and prawns and nutritious ones for the plants in the form of nitrates. The cleanest and most nutritious water was transferred to the channel with a substrate bed (limestone) used as a hydroponic system where the lettuces were planted. This water was then collected in a tank and pumped back to the aquaculture tanks to close the recirculation system.

The greenhouse aquaponic system, with a plastic roof and fine anti-aphid mesh walls, maintained on average a relative humidity of 65 percent and an ambient temperature of 35 degrees-C during the day. Dissolved oxygen levels were close to saturation (from 6.9 to 10.1 with an average of 8.7 mg/L) during the study period, and water temperature fluctuated between 27.0 and 31.0 degrees-C, with the highest values 32.4 degrees-C measured during the afternoons. All water quality parameters measured were within the values reported as adequate for the species tested.

Table 1 shows the final production results for the 85-day study period for the catfish and prawns

The survival rate for the prawns was 86 percent and for the catfish it was 96 percent, reflecting the appropriate operation of the aquaponic system to support both species. Prawn survival was related to the installation of a mesh that served as a natural refuge.

## Valverde, Aquaponics-Costa Rica, Table 1

Parameter	PW Prawns	Catfish
Initial weight (g)	0.003	0.5
Number stocked	200	100
Culture period (days)	85	85
Number harvested	172	96

Survival (%)	86	96
Mean final weight (g)	1.8 ± 1.5	9.9 ± 4.2
Growth (g/week)	0.15	0.82
Biomass (g)	309.6	950
Feed amount (g)	324	1,600
Feed rate (g/m <sup>3</sup> x day)	3.8	18.8
FCR	1.0	1.7

Table 1. Production data for the catfish and prawns cultured during the study.

The final average weights after 85 days of culture was 9.9 grams for the catfish and 1.8 grams for the prawns. Both can be considered low due to the high stocking densities used. In the case of prawns, it could represent a high-density nursery stage for juveniles – resulting in higher survivals because of the controlled culture environment – that can later be transferred to other culture systems and stocked at lower densities for grow-out.

Fig. 1: Size distribution of the two prawn populations that developed in the study.

Fig. 1 shows prawn size distribution, and the normal distribution of two different populations is evident. The largest prawns are those that were stocked first and had final weights from 2 grams to over 5 grams. The small prawns are from the population stocked a month later and had final weights from less than 0.5 grams and up to 2 grams. The size disparity increased as the prawns grow larger.

Representative prawns from the smallest to the largest sizes produced, and channel catfish (r).

The growth rates were 0.15 and 0.82 grams per week for prawns and catfish, respectively. In pond culture, these species reach an average of 20 grams for prawns and 600 grams for catfish after 7 months (210 days) of culture. This means that the growth is 0.67 grams/week and 20.0 grams/week respectively. Therefore, we consider the growth rate for both species extremely low in our aquaponics system.

The biomasses reached during the growing period were 310 grams and 950 grams for the prawns and the catfish, respectively. During the study period, the prawns were fed to satiety with 324 grams and the catfish were fed 1,600 grams of feed. The resulting feed conversion were 1:1 and 1.7:1, respectively; both of these values are within standards for pond culture of these species, so feeding was adequate in our experimental system.

## Valverde, Aquaponics-Costa Rica, Table 2

Parameter	Prawn tank	Catfish tank	Aquaculture tank drainage	Plant drainage
Iron (mg/L)	0.00 ± 0.00	0.03 ± 0.02	0.00 ± 0.00	0.04 ± 0.06
Nitrate (mg/L)	2.75 ± 0.21	4.60 ± 0.99	4.45 ± 2.05	4.00 ± 1.98
Nitrite (mg/L)	0.05 ± 0.00	0.05 ± 0.00	0.04 ± 0.01	0.05 ± 0.00
TAN (mg/L)	0.00 ± 0.00	0.06 ± 0.06	0.03 ± 0.01	0.05 ± 0.06
Chlorides (mg/L)	0.09 ± 0.04	0.12 ± 0.02	0.08 ± 0.00	0.09 ± 0.00
Alkalinity (mg/L CaCO <sub>3</sub> )	110 ± 7	112 ± 9	110 ± 2	117 ± 25
Hardness (mg/L Ca CO <sub>3</sub> )	125 ± 15	174 ± 53	142 ± 6	165 ± 11

Table 2. Various water quality parameters determined at different locations of the experimental aquaponics system.

Fig. 2: Foliage and root lengths for the lettuce varieties studied.

Fig. 2 shows the yields of the three lettuce varieties in terms of foliage and root lengths. Our results show significantly lower performance in these variables for the Rose variety versus the American and Romaine varieties. Of the number of leaves counted in this variety, 30 percent were dead or very deteriorated. The Romaine variety had significantly greater root development than the American variety, while there were no significant differences in foliage height or number of leaves between these varieties.

*References available from first author.*

## Authors

---



### **JOSÉ VALVERDE**

Núcleo Náutico Pesquero  
Instituto Nacional de Aprendizaje (INA)  
Puntarenas, Costa Rica

[jvalmo@yahoo.com](mailto:jvalmo@yahoo.com) (<mailto:jvalmo@yahoo.com>).



### **MARCO ACOSTA**

Núcleo Náutico Pesquero  
Instituto Nacional de Aprendizaje (INA)  
Puntarenas, Costa Rica



**MARVIN TORRES**

Dirección de Investigaciones  
Sede Central, UTN  
Costa Rica

Copyright © 2023 Global Seafood Alliance

All rights reserved.