



ALLIANCE™


<https://debug.globalseafood.org>Health &
Welfare

Ich parasite serves as vector to transmit bacteria to fish

2 November 2012

By De-Hai Xu, Ph.D. , Craig Shoemaker, Ph.D. and Phillip Klesius, Ph.D.

Study evaluates the vector function of parasite for *Edwardsiella ictaluri*



The control theronts (left) showed no bacteria, but many theronts exposed to *E. ictaluri* (right) carried fluorescent bacteria.

Parasites can cause fish death or poor health by directly damaging organs, utilizing nutrients and thus impacting growth, causing stress and thus reducing immune protection against pathogens, and damaging fish epithelium in skin and gills, aiding bacterial entry into the fish.

Fish mucus covers the epithelium and provides natural protection against pathogens. Studies have demonstrated that parasite injuries act as potential portals of entry for bacterial pathogens. Parasitic infection can enhance bacterial invasion and result in high fish mortality.

There is limited information available on whether parasites act as vectors to transmit pathogenic bacteria to fish. Therefore, a study was conducted to evaluate the vector function of the parasite *Ichthyophthirius multifiliis* (ich) for the bacterium *Edwardsiella ictaluri* into channel catfish. Protozoan ich and *E. ictaluri* are common pathogens of cultured channel fish that can result in high mortality and decreased profits. The life cycle of ich consists of an infective theront, a parasitic trophont and a reproductive tomont.

***E. ictaluri* attachment**

The authors evaluated whether the bacteria could attach to ich theronts using green fluorescent-tagged *E. ictaluri*. Nine tubes were filled with theront solution, and the tagged *E. ictaluri* were added at concentrations of 0 colony-forming unit, 4×10^3 CFU and 4×10^7 CFU/mL.

The theronts were exposed to *E. ictaluri* for one hour in the tubes used for each *E. ictaluri* concentration. At the end of the exposure, 1 percent formalin was added to each tube to fix the theronts for 30 minutes.

Theronts were harvested after washing three times with sterile water, then suspended in 0.5 ml of sterile water in flow cytometer tubes. The number of theronts carrying fluorescent *E. ictaluri* was counted for each sample using a flow cytometer. Theronts without *E. ictaluri* exposure were included as negative controls.

The control theronts showed no bacteria, but many theronts exposed to *E. ictaluri* were shown to carry fluorescent bacteria. Theronts (24 and 40 percent) showed fluorescent bacteria after exposure to *E. ictaluri* at 4×10^3 and 4×10^7 CFU/mL, respectively (Table 1).

Hai Xu, Numbers of ice throngs, Table 1

Concentration of <i>E. ictaluri</i>	Theronts Counted	Fluorescent Theronts*	Theronts Positive for <i>E. ictaluri</i>
0 CFU/mL	931	53	5.7%
4×10^3 CFU/mL	992	233	23.5%
4×10^7 CFU/mL	956	321	39.0%

* Theronts showed weak auto-fluorescence.

Table 1. Numbers of ich theronts positive for *E. ictaluri* one hour after exposure.

Tomont exposure

To evaluate whether tomonts exposed to *E. ictaluri* produced theronts carrying *E. ictaluri*, the authors added 300 tomonts per well to duplicate six-well plates. *E. ictaluri* were added to the wells at concentrations of 0, 4×10^5 and 4×10^7 CFU/mL for a two-hour exposure.

The bacterial suspension and unattached tomonts were removed from each well. Then fresh water was added to each well to wash the attached tomonts and remove suspended bacteria. After washing, 30 mL of fresh tank water was added to each well and incubated at 22 ± 2 degrees C. One plate was sampled four or 24 hours after exposure to *E. ictaluri*. The attached tomonts (four hours) or theronts (24 hours) were harvested and viewed with a fluorescent microscope.

No fluorescent bacteria were observed on the control tomonts. All tomonts demonstrated fluorescent bacteria four hours post-exposure to *E. ictaluri* (Table 2). After 24 hours, most tomonts divided into several hundred infective theronts. Among those theronts, 31 percent and 66 percent were observed to have bacteria attached to their surfaces following tomont exposure to *E. ictaluri* at 5×10^5 or 5×10^7 CFU/mL (Table 2). *E. ictaluri* survived and replicated during the tomont division.

Hai Xu, Number for torments, Table 2

Concentration of <i>E. ictaluri</i>	Hour 4 P/N	Hour 4 % P	Hour 24 P/N	Hour 24 % P
0 CFU/mL	0/31	0	0/261	0
5×10^5 CFU/mL	45/45	100		31.2
5×10^7 CFU/mL	40/40	100	77/247	66.4

P = Number of tomonts or theronts positive for *E. ictaluri*

N = Number of tomonts or theronts examined

% P = Percentage of tomonts or theronts positive for *E. ictaluri*

Table 2. Number of tomonts and theronts positive for *E. ictaluri*.

Fish infection trial

E. ictaluri were added to ich theront solution in three, 1-L beakers at concentrations of 0, 4×10^5 and 4×10^7 CFU/mL. After exposure to *E. ictaluri* for one hour, theronts were harvested and washed. Six, 2-L beakers each were filled with 1 L of water and 30 channel catfish fingerlings weighing 0.3 ± 0.1 g. The theronts exposed to various concentrations of *E. ictaluri* were added to each beaker at 1,000 theronts/fish with two beakers for each treatment. Five fish were sampled from each beaker four hours and one day after theront exposure. Tissue from each sample was grown in bacterial medium for 24 hours to examine the presence of *E. ictaluri*.

Approximately 60 percent and 90 percent of the fish exposed to theronts treated with 5×10^5 *E. ictaluri*/mL showed fluorescent bacteria at four hours and one day (Table 3). All fish were positive for *E. ictaluri* four hours after exposure to theronts treated with 5×10^7 *E. ictaluri*/mL. Two days post-

exposure, cumulative fish mortalities were 36.7 percent, 40.0 percent, and 60.0 percent for fish exposed to theronts only, theronts treated with 5×10^5 *E. ictaluri*/mL and theronts treated with 5×10^7 *E. ictaluri*/mL, respectively.

Hai Xu, Number of fish positive, Table 3

Concentration of <i>E. ictaluri</i>	Hour 4 P/N	Hour 4 % P	Hour 24 P/N	Hour 24 % P
0 CFU/mL	0/10	0	0/10	0
5×10^5 CFU/mL	6/10	60	9/10	90
5×10^7 CFU/mL	10/10	100	10/10	100

P = Number of fish positive for *E. ictaluri*

N = Number of fish examined


% P = Percentage of fish positive for *E. ictaluri*

Table 3. Number of fish positive for *E. ictaluri* after exposure to theronts treated with *E. ictaluri*.

Results of this study demonstrated that ich can vector *E. ictaluri* into channel catfish. Understanding the potential ability of parasites to vector bacterial disease is important to fish farmers and health managers particularly because parasites introduced via wild fish or fish from other farms could concomitantly involve the introduction and/or transmission of microbial disease agents.

(Editor's Note: This article was originally published in the November/December 2012 print edition of the Global Aquaculture Advocate.)

Authors



DE-HAI XU, PH.D.

USDA Agricultural Research Service
Aquatic Animal Health Research Laboratory
990 Wire Road
Auburn, Alabama 36832 USA

dehai.xu@ars.usda.gov (mailto:dehai.xu@ars.usda.gov)



CRAIG SHOEMAKER, PH.D.

USDA Agricultural Research Service
Aquatic Animal Health Research Laboratory
990 Wire Road
Auburn, Alabama 36832 USA



PHILLIP KLESIUS, PH.D.

USDA Agricultural Research Service
Aquatic Animal Health Research Laboratory
990 Wire Road
Auburn, Alabama 36832 USA

Copyright © 2023 Global Seafood Alliance

All rights reserved.