



ALLIANCE™

(<https://debug.globalseafood.org>).

---

Intelligence

---

# Pearl oyster culture in Mexico

1 June 2003

By Héctor Acosta-Salmón

## Potential for farms along the Gulf of California, Mexican Pacific coast



The U.S. \$16,000 "Isabella" necklace has 43 round Mexican pearls.


Photo by E. Arizmendi, courtesy of Perlas del Mar de Cortez.

Pearl culture is one of the more valuable aquaculture industries in the world, with an estimated retail value of around U.S. \$5 billion per year. Most of the world's marine pearls are produced by western Pacific pearl oysters (*Pinctada maxima*, silver or gold pearls), *P. margaritifera* (black pearls) and *P. fucata* (white pearls). Japan, Australia and French Polynesia are the main pearl-producing countries.

Scientific research has focused mainly on the three pearl oyster species above. The past 15 years have provided the most significant advances in hatchery spat production, genetic manipulation and pre- and postoperative grafting treatments.

## Pearl oyster production in Mexico

Mexico was a pioneer in culturing pearl oysters. Before the end of the 19<sup>th</sup> century, Compañía Criadora de Concha y Perla, a large venture based in La Paz Bay, collected wild pearl oyster spat and grew the oysters to adult stage. Seeding operations were not performed, and only natural pearls and shells were harvested. The company held in culture over 3 million adult pearl oysters. In the early 1900s, the company was destroyed during the Mexican Revolution and never recovered.



Adult *P. mazatlanica* cultured in plastic cages. Photo by A. Olivera.

Afterwards, pearl oyster fishing fleets increased in numbers. But after an unusually high mortality attributed to such causes as overfishing, El Niño, the closing of the Colorado River as it flowed into the Gulf of California, and even deliberate poisoning, the federal government decreed a total ban on fishing for pearl oysters in 1939. Nowadays, wild populations have still not recovered, and illegal fishing activities remain a problem.

During the 1960s, attempts to establish a pearl farm in La Paz Bay were abandoned when the Mexican government stopped the project due to lack of appropriate legislation and disagreements between the authorities and farm management. This lack of support prevented any further new projects.

At around this time, the pearl industry of French Polynesia began to develop. Forty years later, Tahitian black pearls are well-known, and Mexican pearling is still in its infancy.

## Species

Two pearl oyster species, *Pinctada mazatlanica* and *Pteria sterna*, are found on the Pacific coast of Mexico and in the Gulf of California. *P. mazatlanica* are very similar in appearance to the black-lip pearl oyster *P. margaritifera*. *Pt. sterna* are smaller than their counterpart, *Pt. penguin*.

*P. mazatlanica* show a distinctive metallic-green lip with overtones in blue or pink. Their pearls range in color from white to dark gray with blue and green hues, and can reach 13-15 mm in diameter. *Pt. sterna* show probably the largest range of nacre colors of any pearl oyster species: pale white, blue, green, purple, pink, gray, and gold. These pearls attain an average diameter of 9 mm. International experts have recognized the quality of the nacre and pearls of these two species of Mexican pearl oysters.

## Research



Bacoachibampo Bay is home to Perlas del Mar de Cortez. La Paz Bay is where most of the scientific research in pearl oysters has been developed.

Early studies were focused on reestablishing natural population stocks or creating new populations through the transplanting of oysters to selected places showing low abundance. Success was very limited until the early 1990s, when the basics for repopulation were established.

During the last 10 years, pearl oyster research in Mexico increased significantly in areas such as larval rearing and settlement in the hatchery, wild spat collection, nursery culture techniques, repopulation, reproductive biology, and pearl production of “mabe,” or half pearls and round pearls. Research

institutions such as Centro de Investigaciones Biológicas del Noroeste, Instituto Tecnológico y de Estudios Superiores de Monterrey (ITESM), and Universidad Autónoma de Baja California Sur have received national, international, and institutional funds to develop pearl oyster culture technology.

Thanks to these projects, some groups have started producing mabe on a commercial scale, and Mexican pearls are now seen in national and international markets. Also, repopulated banks of pearl oysters have been successfully maintained in La Paz Bay.

## Industry

In 1994, the first cultured half pearls were obtained under experimental and pilot scale, but it was not until 1996 when the first commercial harvest of mabe from *Pt. sterna* was obtained by ITESM in Bacochibampo Bay, Guaymas. ITESM production is currently around 5,000 mabe and 5,000 round pearls per year. The unique necklace "Isabella," with 43 round 8.0- to 10.5-mm Mexican pearls, was sold for U.S. \$16,000 last year.

During the past several years, small groups have started to collect wild spat to culture *Pt. sterna* and produce pearls for the local markets. Plans to develop a pearl quality control policy are under way by these groups. People from within each group perform all nuclei grafting operations.

Though hatchery production is a reality today, it is limited to research activities. Enough wild spat, especially of *Pt. sterna*, can be collected for the current small commercial activities.

## Conclusion

The characteristics of Mexican pearls make them a unique product with limited competition in the international market, but great care must be taken when offering them to the world. Strict quality control policies must be implemented to assure a minimum standard of quality and protect consumers as well as the incipient Mexican pearl industry.

Scientific research will continue to focus on reproductive biology as well as pearl production, with a special emphasis on pearl quality. As all the groups have evolved from teaching research institutions, they will also continue scientific research.

There is potential for establishing pearl farms along the Gulf of California and the Mexican Pacific coast. The most likely scenario is that commercial pearl farms will evolve from the small groups currently producing mabe. Other aquaculture groups, such as those working with edible oysters or marine fish, could also start collecting wild pearl oyster spat to diversify their operations and participate in community-based pearl oyster culture programs. The production of round pearls will improve slowly within each group, as the cost to hire expert seeding technicians cannot currently be met.

Another approach to facilitate development of the pearl industry in Mexico would be for state governments to finance local fishing communities to start cooperative pearl farms. Finally, selective breeding and stock enhancement may be the trigger for commercial farms to give more attention to hatchery-produced spat.

*(Editor's Note: This article was originally published in the June 2003 print edition of the Global Aquaculture Advocate.)*

## Author

---



**HÉCTOR ACOSTA-SALMÓN**

Pearl Oyster Research Group  
School of Marine Biology and Aquaculture  
James Cook University  
Townsville, Queensland 4811  
Australia

[hector.acostasalmon@jcu.edu.au](mailto:hector.acostasalmon@jcu.edu.au) (<mailto:hector.acostasalmon@jcu.edu.au>).

Copyright © 2023 Global Seafood Alliance

All rights reserved.