




ALLIANCE™

(<https://debug.globalseafood.org>).

---



Health &  
Welfare


---

# Real-time PCR offers sensitivity, specificity in detecting AHPND plasmid

2 May 2015

By Jee Eun Han, DVM, Ph.D. , Kathy Tang, Ph.D. , Brenda White and Donald Lightner, Ph.D.

## Quantification of pathogen in infected animals crucial for monitoring disease progression



Highly sensitive new assays can be used for quantification of pathogens in water and shrimp samples from farms.

Acute hepatopancreatic necrosis disease (AHPND), also known as early mortality syndrome, has caused severe mortalities in farmed populations of *Litopenaeus vannamei* and *Penaeus monodon* shrimp. The disease has led to significant production and economic losses at shrimp farms and to the aquaculture industry in general in affected regions.

Clinical signs of the disease in infected shrimp include an empty gastrointestinal tract and whitish, atrophied stomach and hepatopancreas. The causative agent was determined to be the unique strains of the bacterium *Vibrio parahaemolyticus* that contain the *pirA*- and *pirB*-like toxin genes. These genes are located in a large, 69-kb plasmid. Plasmids are self-replicating, double-stranded DNA molecules that pass between bacterial cells through conjugation and transformation.

Initially, diagnosis of AHPND could only be accomplished through histological examinations and laboratory bioassays. Later, molecular diagnostic methods based on conventional polymerase chain reaction testing targeting *pirA*- and/or *pirB*-like genes became available. However, quantification of the AHPND pathogen in the infected animals is one of the most important means for monitoring disease progression.

Bacterial quantities can be determined through counting colony-forming units (CFUs), but this is a laborious and time-consuming method. Real-time, quantitative polymerase chain reaction (qPCR) analysis has become an attractive alternative for its advantages of speed, sensitivity and specificity.

## PCR primers, probe

The PCR primers and specificity-enhancing hydrolysis probe for the detection of the virulence plasmid were selected from the *pirA*-like gene. The primers were used to amplify a DNA fragment of 135-bp. The probe was synthesized and labeled with different dyes on either end.

Samples of extracted DNA were added to a qPCR mixture containing 0.3  $\mu$  of each primer and 0.1  $\mu$  probe to a final volume of 10  $\mu$ L. The qPCR profile consisted of 20 seconds at 95 degrees C, followed by 40 cycles of 3 seconds at 95 degrees C and 30 seconds at 60 degrees C.

## Sensitivity, specificity

The detection limit of this qPCR assay was less than 10 copies of virulence plasmid. The standard curve ( $10^2$  to  $10^8$  copies of plasmid) is shown in Figure 1. This method can detect 12 AHPND-causing *V. parahaemolyticus* isolates from Mexico and Vietnam, but did not cross react with 35 non-pathogenic *Vibrio* bacteria.




Fig. 1: Standard curve and gel electrophoresis of the AHPND qPCR assay. Insert: Gel electrophoresis of amplified products from qPCR.

## Affect on farmed shrimp

The authors used two cases of farm-raised *L. vannamei* collected in 2012 to detect and quantify the AHPND plasmid. Case 1 consisted of 32 small juvenile shrimp collected from China that showed significant mortalities and severe effects of AHPND, determined by histological examination.

By qPCR, DNA extracted from each shrimp proved to be AHPND-positive. The quantities of virulence plasmid ranged from  $2.5 \times 10^3$  to  $4.7 \times 10^6$  copies per milligram of tissue (Table 1).

Laboratory Bioassay	Infection Route	Shrimp	Copies/mg Tissue
1	Per os (1 x 10 <sup>9</sup> CFU/ml)	Day 1 (moribund/dead) Day 2 (moribund/dead)	2.5 x 10 <sup>3</sup> -4.7 x 10 <sup>6</sup> 2.1 x 10 <sup>5</sup> -3.6 x 10 <sup>6</sup>
2	Immersion (2 x 10 <sup>5</sup> CFU/ml)	Day 1 (moribund/dead) Day 2 (moribund/dead) Day 9 (surviving)	1.8 x 10 <sup>3</sup> -2.1 x 10 <sup>5</sup> 1.4 x 10 <sup>5</sup> -7.9 x 10 <sup>5</sup>
3	Per os (8 x 10 <sup>8</sup> CFU/mL)		1.8 x 10 <sup>1</sup> -3.9 x 10 <sup>2</sup>
4	Per os (1 x 10 <sup>9</sup> CFU/mL)	Water sample	3.5 x 10 <sup>2</sup> -2.2 x 10 <sup>6</sup> *

The second case included 12 juvenile shrimp collected from the Ben Tre Province of Vietnam. These shrimp suffered high mortalities and were later found to have AHPND through histological examination. In qPCR analysis, DNA extracted from hepatopancreas tissues found five of the 12 shrimp were positive for AHPND. The quantities of virulence plasmid were 3.9 to 5.8 x 10<sup>5</sup> copies per milligram of tissue.

Virulence plasmid copy numbers varied widely, ranging 4-10<sup>5-6</sup>/mg tissue. Thus, the AHPND bacterial load in affected shrimp would be expected to also be highly variable. This suggested that factors other than bacterial load, such as shrimp age or size, the presence of other pathogens or environmental conditions, may influence the mortalities in affected farms.

## Laboratory-infected shrimp

Shrimp collected from laboratory infections were analyzed for AHPND. In bioassay 1, a per os feeding experiment, specific-pathogen-free *L. vannamei* with a mean individual weight of 1 g were fed shrimp feed mixed with AHPND *V. parahaemolyticus*. The bacteria were first grown to 1 x 10<sup>9</sup> CFU/mL and mixed with shrimp feed at a 1:1 ratio.

All 50 infected shrimp became moribund or died within two days. These shrimp were positive for AHPND by histological examination and qPCR. The quantities of virulence plasmid ranged from 2.5 x 10<sup>3</sup> to 4.7 x 10<sup>6</sup> per milligram of tissue (Table 1).

In bioassay 2, shrimp were exposed to AHPND *V. parahaemolyticus* by immersion at a concentration of 2 x 10<sup>5</sup> CFU/mL water. All the shrimp died within two days. By qPCR, the quantities of virulence plasmid in the moribund/dead shrimp ranged from 1.8 x 10<sup>3</sup> to 7.9 x 10<sup>5</sup> copies per milligram of tissue (Table 1).

Bioassay 3 was a challenge study to a selected line of *L. vannamei*. At a mean weight of 2 g, the shrimp were stocked in 1,000-L tanks and given feed containing AHPND *V. parahaemolyticus*. At day 9, 244 shrimp (48 percent) survived, whereas none of the 50 shrimp in the positive control tank survived after two days.

Histological examination did not detect AHPND in these surviving shrimp. By qPCR, the quantities of AHPND plasmid were 18-390 copies/mg tissue (Table 1), suggesting this selected line was better at fending off the disease. However, these surviving shrimp, which carried low quantities of pathogenic bacteria, could act as vectors capable of spreading the disease.


Although hepatopancreas tissue is usually sampled for detecting AHPND *V. parahaemolyticus*, a fourth bioassay was performed to demonstrate that water samples can be used for AHPND diagnostics.

In this bioassay with shrimp fed feed mixed with AHPND *V. parahaemolyticus*, water was sampled from each tank at day 5 and concentrated 50 times. AHPND was detected in seven of nine water samples by the highly sensitive qPCR, ranging  $3.5 \times 10^2$  to  $2.2 \times 10^6$  plasmid copies/mL water (Table 1). This is relevant, since it has been reported that AHPND can be transmitted via water from AHPND-affected ponds.

*(Editor's Note: This article was originally published in the May/June 2015 print edition of the Global Aquaculture Advocate.)*

## Authors

---



**JEE EUN HAN, DVM, PH.D.**

School of Animal and Comparative Biomedical Sciences  
University of Arizona  
Tucson, Arizona 85721 USA

[jeehan@email.arizona.edu](mailto:jeehan@email.arizona.edu) (<mailto:jeehan@email.arizona.edu>)



**KATHY TANG, PH.D.**

School of Animal and Comparative Biomedical Sciences  
University of Arizona



**BRENDA WHITE**

School of Animal and Comparative Biomedical Sciences  
University of Arizona



**DONALD LIGHTNER, PH.D.**

School of Animal and Comparative Biomedical Sciences  
University of Arizona

Copyright © 2023 Global Seafood Alliance

All rights reserved.