




ALLIANCE™

(<https://debug.globalseafood.org>).



 Health & Welfare

# Surveillance an important health management tool

1 March 2009

By Victoria Alday-Sanz, Ph.D.

Protect health with **water filtration, pond netting and use of domesticated stocks**



Regular monitoring and sampling of aquaculture ponds can reveal animal condition and subsequent disease problems before they get out of hand.

Regular surveillance provides information on the health status of farmed stocks. It can also provide information on possible future disease outbreaks by detecting increases in the prevalence of endemic pathogens or the presence of exotic pathogens. Their early detection can allow intervention and control measures before disease outbreaks occur.

### **Early detection**

Surveillance can be focused on known or unknown pathogens, or a combination of both. For known pathogens, polymerase chain reaction techniques should be applied. Each farm should determine its own list of important diseases to protect against. In addition to other criteria, this list could be based on the economic impacts of the diseases, means of transmission and presence of susceptible species.

For unknown pathogens, histology is required. Histological changes, even if the causative agent cannot be identified, should raise the attention of the farmer. If there is an increase in the prevalence of these changes, they may reflect the build-up of an epidemic.

### **Improving performance**

The line that separates healthy and sick animals in aquaculture is rather blurred. From the point of view of a farmer, health is a measure of productivity. If a disease does not cause unexpected loss of income, it may not be regarded as a disease concern. However, chronic diseases that reduce growth and survival have a significant cost to the industry.

Often, results of diagnostic tests are used for immediate application of management practices, such as stocking or discharge of postlarvae, emergency harvest or exchange of water. However, this information can also be used for medium-term management. Some epidemiological studies indicate that a random whole farm sampling for all pathogens could be more advantageous than a thorough follow-up on a small number of ponds.

When surveillance detects pathogens such as hepatopancreatic parvovirus in Thailand or infectious hypodermal and hematopoietic necrosis virus in Latin America, or risk factors associated with low productivity, a targeted sampling could be implemented as a monitoring program in following cycles. Such programs could optimize the value of the diagnostic information and identify ways to improve production.

### **Biosecurity needed**

In an interesting paper on the emergence of viruses in Australian prawn aquaculture published in 1997 in the *World Journal of Microbiology & Biotechnology*, Dr. Leigh Owens of James Cook University in Australia presented the detection of a number of viruses in wild shrimp, wild-caught broodstock and farmed shrimp within a time frame (Table 1).

## **Alday-Sanz, Dates of detection of shrimp viruses, Table 1**

<b>Virus, Condition</b>	<b>Wild Shrimp</b>	<b>Wild Broodstock</b>	<b>Aquaculture</b>
RNA proliferative syndrome	1980		
Hepatopancreatic parvovirus	1984		1988
Monodon baculovirus	1984	1986	1986
Lymphoid organ vacuolization	1988		1990
Lymphoidal parvovirus	1989		1990
Infectious hypodermal and hematopoietic necrosis virus	1989	1991	
Penaeus hybrid rod-shaped virus	1989	1991	1996
Gut and nerve syndrome	1993		
Spawner-isolated mortality virus		1993	1994
Lymphoid organ virus		1995	1995
Bennettiae baculovirus	1995		
Midgut caecum inclusions			1990

Table 1. Dates of detection of shrimp viruses in Australia, Owens 1997.

The paper showed that early in Australia's aquaculture history, the detection of all viruses was preceded by their discovery in wild prawns from the fisheries, highlighting the risk of introducing wild broodstock to farm facilities. Once pathogens are introduced to farms, they find an ideal environment for replication, spread and potentially lead to disease outbreaks.

The health status of culture species can therefore be protected through improvements in water filtration, pond netting and the use of only domesticated stocks. While new viruses will be discovered in the future, only by closing the life cycle and improving biosecurity can we start excluding pathogens from aquaculture production in the present.

*(Editor's Note: This article was originally published in the March/April 2009 print edition of the Global Aquaculture Advocate.)*

## Author



**VICTORIA ALDAY-SANZ, PH.D.**

Aquatic Animal Health  
Gran Via 658, 4-1  
08010 Barcelona, Spain

[victoria\\_alday@yahoo.com](mailto:victoria_alday@yahoo.com) (mailto:victoria\_alday@yahoo.com)

Copyright © 2023 Global Seafood Alliance

All rights reserved.